
Unusual Lateral Presentation of Popliteal Cyst

Tarek Hemmali,*

Abstract:

The most common cyst occurs in the popliteal region is the popliteal cyst and over the past years it has been received much clinical attention. The most common position of the popliteal cyst is in the posterosuperior aspect of the medial femoral condyle. The present paper describes a rare case of popliteal cyst, which was found in the postero-superior aspect of the lateral condyle of the femur. Interestingly, there are fewer reports of popliteal cysts occurring in the postero-superior aspect of the lateral condyle. We as orthopedic surgeons describe the unusual presentation of popliteal cyst in the postero-superior aspect of the lateral popliteal region and discuss its clinical significance. Anatomical knowledge of such anomalies is very important.

Introduction:

Popliteal cyst, also known as Baker's cyst is the most commonly occurring mass in the popliteal region. The classical Baker's cyst presents as an enlargement of the gastrocnemius-semimembranosus bursa and is related to the postero-superior aspect of medial femoral condyle. Later presentation of the cyst in the popliteal region is rare. The cyst that is being described in the present case was related to lateral femoral condyle and extended proximally on the popliteal surface of femur deep to the popliteal vessels. Such a presentation may be confused clinically with common swellings in this area such as

thrombophlebitis, lipoma or xanthoma. The anatomical knowledge of such rare variations might be helpful for the diagnosis and careful clinical assessment of the popliteal swellings prior to any surgical procedure.

Case Presentation:

A 37 years old Libyan male came to us and complaining about swelling in popliteal fossa left knee for 6 wks.

It was painless swelling.

On examination there was a 5×8 cm mass occupying lateral aspect of popliteal fossa, soft in consistency.




*) Department of Orthopedic Surgery, Tripoli Central Hospital, Tripoli, Libya.










MRI Report:

Mild joint effusion with grade 1 degenerative tear in the posterior horn of the medial meniscus.
A Large popliteal cyst at postero-lateral aspect of the lateral femoral condyle of the knee joint, lateral to lateral head of gastrocnemius.
It measures about 4.2x5.5x7cm (AnteroPosterior x width x height respectively)












Operation Note:


Large lobulated tense swelling 8x5x3cm.
The cyst was related medially to popliteal blood vessels and tibial nerve and laterally to the biceps femoris muscle tendon.
It was closely associated to the popliteal surface of the femur. On careful observation, the cyst was found to be translucent, fluctuant, containing viscous reddish fluid and adherent to adjacent structures.

The cyst was present on the postero-superior aspect of lateral condyle of femur and extended proximally and distally on its popliteal surface. Further exploration revealed a communication between the cyst stalk and the joint cavity. No abnormality as regards the position of the blood vessels and nerves was noted in this region

The cyst was excised; the cyst stalk ligated and sent for histopathological examination.









Discussion:

Baker's cyst is a synovial cyst in the popliteal fossa formed either by the accumulation of synovial fluid in a non-communicating bursa, distension of the bursa by the synovial fluid from the knee joint or by the herniation of the posterior part of capsule due to increased intra-articular pressure.¹ It is characteristically located at the postero-medial aspect of knee joint and extends distally into the calf.² It is frequently associated with degenerative joint disease or may occur spontaneously without any intra-articular pathology.³ The lateral presentation of Baker's cyst was relatively uncommon.⁴⁻⁶

In the present report, the cyst was related to the posterior aspect of lateral condyle of femur and extended proximally in the region of thigh. Another rare feature in the present study was that this cyst was present deep to popliteal vessels and tibial nerve while in the earlier reported cases it was superficial to them.¹⁻³ A thorough review of literature revealed only a few reports of lateral presentation of popliteal cysts.⁴⁻⁶ Only a single case of anterior presentation of such a cyst and another of lateral Baker's cyst herniating through the iliotibial tract have also been reported.^{5,7} An intramuscular cyst located within the vastus lateralis in MRI studies have been reported.⁸

The popliteal cyst in the present case arose laterally and communicated with the joint cavity. Such cysts on the lateral side of knee joint may be bursal or synovial in origin. The

synovial membrane may not give an anatomical support thus giving rise to the cyst.¹⁻³ Since there was no evidence of any meniscal tear, the chances of a meniscal cyst can be ruled out.

When distended, one of the laterally placed bursae, especially that between the lateral femoral condyle and the tendons of popliteus or biceps femoris may present as a lateral cyst.⁴⁻⁶ Interestingly a past study had detected a lateral cyst only in a CT scan.⁷

Due to its unusual presentation, it may mimic aneurysm, deep vein thrombosis, thrombophlebitis, hematoma, xanthoma, abscess or soft tissue tumour of the calf.⁹ A proximate position of the cyst to the deep veins and neuro vascular structures also increases the chances of the involvement of the latter.

It may be presumed that this cyst could have arisen from an enlarged communicating bursa or possibly herniation of the synovial membrane through the posterior part of capsule. Admittedly, we could not carry out the histological study to arrive at a certain diagnosis.

The anatomical knowledge of such cysts is also important during ultrasonographic studies.¹⁰

This paper attempts to highlight the unusual lateral position of popliteal cyst, the anatomical knowledge of which would be significant for clinicians and orthopedic surgeons.

References:

1. Meyerding HW, VanDemark RE. Posterior hernia of the knee (Baker's cyst, Popliteal cyst, Semimembranosus Bursitis, Medial Gastrocnemius Bursitis and Popliteal Bursitis). *J Amer Med Assoc* 1943; 122: 858.
2. Labropoulos N, Shifrin DA, Paxinos O. New insights into the development of popliteal cysts. *Brit J Surg* 2004; 91: 1313-8.
3. Handy JR. Popliteal cysts in adults: a review. *Semin Arthritis Rheum* 2001; 31: 108-18.
4. Buckwalter JA, Dryer RF, Mickelson MD. Arthrography in juxta-articular cysts of the knee. *J Bone Joint Surg* 1979; 61A: 465.
5. O'Dell JR, Andersen PA, Hollister JR, West SC.). Anterior tibial mass: An unusual complication of popliteal cysts. *Arthritis Rheumatis* 1984; 27: 113.
6. Perri JA, Rodman GP, Mankin HJ. Giant synovial cysts of the calf in patients with rheumatoid arthritis. *J Bone Joint Surg* 1968; 50-A: 709-19.
7. Jensen KH, Jorgensen U. Lateral presentation of a Baker's cyst. *Clinical Orthopaedics* 1993; 287: 202-203.
8. Burk DL, Dalinka MK, Kanal E et al. Meniscal and ganglion cysts of the knee. *Amer J Roentgenol* 1988;150: 331-6.
9. Schmidt MC, Workman JB, Bartu WF. Dissection or rupture of a popliteal cyst: a syndrome mimicking thrombophlebitis in rheumatic diseases. *Arch Intern Med* 1974; 134: 694-8.
10. Rupp S, Seil R, Jochum P, Kohn D. Popliteal cysts in adults. Prevalence, associated intraarticular lesions, and results after arthroscopic treatment. *Amer J Sports Med* 2002; 30: 112-5.