



The combined effect of biochar and organic fertilizer (goat manure) on the growth and yield of peanut crop (*Arachis hypogaea L.*)

*Abdouslam M Rassem, Khalid A Elzobair

Department of Soil and Water, Faculty of Agriculture, Sabha University, P. O. Box 18758, Sabha, Libya

Keywords:

Biochar
Goat manure
Peanut yield
Plant growth
Sandy soil

ABSTRACT

Sandy soils suffer from many problems with physical and chemical properties as well as low productivity of cultivated crops. Biochar and organic fertilizer have been used in agriculture to improve soil fertility, enhance crop productivity, and sequester carbon in the soil. A study was conducted to investigate the effect of biochar (B) from woody materials and goat manure (GM) on the growth and yield of peanut crops (*Arachis hypogaea L.*) in sandy soil. The experiment was in a randomized complete block design RCBD with three replicates. The treatments were controlled, 10 t ha⁻¹ GM, 2 % B + 10 t ha⁻¹ GM, and 5 % B + 10 t ha⁻¹ GM. The result showed that the addition of 2 % B + 10 t ha⁻¹ GM increased peanut yield by 50 % while the addition of 5 % B + GM 10 t ha⁻¹ increased peanut yield by 82 %. Without biochar, goat manure had yield increases by 25 % compared to a control. Overall, our findings demonstrate that biochar-goat manure combinations have a better performance than GM alone, in terms of yield and plant growth. For future research, we tend to determine the optimal combination of biochar and goat manure for peanut production.

التأثير المشترك للفحم الحيوي والسماذ العضوي (روث الماعز) على نمو وإنتاجية محصول الفول السوداني (*Arachis hypogaeaL*)

*عبد السلام راسم و خالد الزبير

قسم التربة والياه، كلية الزراعة، جامعة سبها، صندوق البريد 18758، ليبيا

الكلمات المفتاحية:

الترب الرملية
الفحم الحيوي
الفول السوداني
روث الماعز
نمو النبات

الملخص

ت تعاني التربة الرملية من مشاكل عديدة في خواصها الفيزيائية والكيميائية بالإضافة إلى انخفاض إنتاجية المحاصيل المزروعة. تم استخدام الفحم الحيوي والأسمدة العضوية في التجربة لتحسين خصوبة التربة، وتعزيز إنتاجية المحاصيل، وحجز الكربون في التربة. أجريت الدراسة لمعرفة تأثير الفحم الحيوي (B) نوع مواد خشبية وروث الماعز (GM) على نمو وإنتاجية محصول الفول السوداني (*Arachis Hypogaea L*) في التربة الرملية. نفذت التجربة بتصميم القطاعات العشوائية الكاملة RCBD بثلاثة مكررات. وأربعة معاملات، الشاهد، 10 طن للهكتار (GM)، 10 % (B) + 2 % طن للهكتار (GM)، و 5 % (B) + 10 % طن للهكتار (GM). أظهرت النتائج أن إضافة 2 % (B) + 10 % طن للهكتار (GM) يزيد إنتاجية الفول السوداني بنسبة 50 % بينما إضافة 5 % (B) + 10 % طن للهكتار (GM) يزيد إنتاجية الفول السوداني بنسبة 82 % بدون الفحم الحيوي وبوجود روث الماعز فقط زاد إنتاج الفول السوداني بنسبة 25 % مقارنة بالشاهد. بشكل عام توضح النتائج التي توصلنا إليها أن إضافة الفحم الحيوي وسماذ الماعز معاً يحسن نمو وإنتاجية محصول الفول السوداني. بالنسبة للأبحاث المستقبلية، فإننا نميل إلى تحديد المزيج الأمثل من الفحم الحيوي وروث الماعز لإنتاج الفول السوداني.

Introduction

Peanut (*Arachis hypogaea L.*) is an annual legume crop that provides food and helps maintain soil fertility through nitrogen fixation [1]

*Corresponding author:

E-mail addresses: abd.rassem@sebhua.edu.ly, (K. Elzobair) elzobair.khalid@gmail.com

Article History : Received 10 June 2023 - Received in revised form 27 September 2023 - Accepted 02 October 2023

The symbiotic relationship between leguminous crops and rhizobium bacteria is becoming increasingly important in the agriculture world, as this potentially leads to more sustainable agricultural systems, reducing requirements for chemical fertilizer, enhancing residual benefits to subsequent crops and increasing crop yields [2]. Globally, the consumption of peanuts is increasing at a rate of around 3 % per annum. In 2011/12, peanut production in the world was 35 million (*United States Department of Agriculture*). China, India, and the USA are the main producers, growing 16.0, 5.5, and 1.7 million tons, respectively, accounting for 45 %, 16 % and 5 % of the world's total respectively [3].

Biochar is a carbon-rich product of burning biomass in the absence of oxygen (pyrolysis), and its potential to improve soil fertility and mitigate climate change has been recognized globally [4][5]. Biochar has a high proportion of recalcitrant C with hundreds to thousands of years of durability, making it a potentially effective soil C sink [6]. Biochar also has unique properties to improve soil productivity [7][8]. The porous physical structure of biochar induces a greater sorption capacity to conserve soil moisture and nutrients. The alkaline nature of many biochars makes such materials especially suitable for improving acidic soil [9]. Biochar made from specific feedstocks (e.g., manure) has high nutrient content and promotes plant growth [10][11][12]. These agronomic benefits have been well demonstrated [13][14][15] especially on marginal and degraded soils [16][15]. However, the fundamental mechanisms by which biochar affects crop growth are insufficiently understood [17]

Recently, biochar has become increasingly the subject of scientific and public interest. It is claimed that biochar can improve soil properties and agronomic performance, inspired by investigations of Terra Preta in Amazon [18]. Several studies showed that the biochar application to soil can influence soil properties (e.g. water holding capacity, pH, and microbial activity) [19][20]. Further studies observed enhanced nutrient uptake by plants after biochar application [14]. In contrast, some authors reported no significant effects of biochar on soil properties, plant nutrition, or biomass production after

biochar application under field conditions [21]. Until now, most biochar studies were performed with pure biochar under laboratory or greenhouse conditions or in tropical environments [15]. Biochar studies under field conditions often show contrasting results to those conducted in the laboratory [22]. Therefore, we conducted a field experiment to quantify the effects of biochar on plant growth when combined with organic fertilizers (goat manure). We amended sandy soils using biochar created from woody material and examined the performance of peanuts on biochar-amended soils in situ experiments.

Material and methods

Location and soil

The field trial was conducted on the farm of Agriculture Facility, Sebha University, Libya. Latitude 26°58'21.58", Longitude 14° 26' 23.85". The experimental land was not cultivated before. The soil is classified as sandy soil (American soil taxonomy; sand 92 %, silt 6.4 %, clay 3.2 %). The mean annual precipitation is 22 mm and the mean annual temperature is 30°C (data source: World Weather Online <https://www.worldweatheronline.com/>). During our study though, the soil site had not received any rainfall.

Experimental design and treatments

The experiment was established in Summer 2022 and was in a randomized complete block design RCBD with three replicates. The treatments were controlled, 10 t ha⁻¹ GM, 2 % B + 10 t ha⁻¹ GM, and 5 % B + 10 t ha⁻¹ GM. Biochar was made in a hole on a micro-scale, ground into a powder of less than 0.5 mm particles, and stored in a sealed plastic bag until use. Biochar at the desired rates (2 or 5 % by weight) was hand applied to the soil surface and till to a depth of 15 cm. Goat manure was obtained from a private farm, and composted for six months under optimum conditions of air and moist content. Other cultural operations were done as per recommendation and crop requirements. The soil samples of the experimental sites were taken at a depth of 30 cm. Some physical and chemical analyses are presented in Table (1): -

Table 1. Chemical and physical properties of soil site.

| Sand | Salt | Clay | pH | EC | O.M | N | P | K |
|------------------------------------|------|------|----------------|--------------------|-------------------------|--------------------------|----------------------------|-----------------------|
| % | | | | dS m ⁻¹ | % | ppm | | |
| 93.65 | 3.64 | 2.80 | 7.2 | 0.48 | 0.44 | 9 | 3 | 80 |
| Sandy | | | Neutral | Very low | Low | Low | low | low |
| Bouyoucos Hydrometer method (1952) | | | Jackson (1958) | Wilcox (1950) | Walkey and Black (1934) | Subbaih and Asija (1956) | Olsen <i>et al.</i> (1954) | Toth and Price (1949) |

The data on morphological, physiological, and yield characteristics were collected. First, we measured plant height (cm), the number of leaves, and the number of branches after 40 and 80 DAS. Then, the number of flowers and the number of pods were taken. Finally, plants were harvested and weighed for the pods, and the biomass.

Statistical analysis

All data were analyzed by statistical analysis software (SAS) version 9.4. To test for significance, we use the analysis of variance (ANOVA). Treatment means were separated for significance using the Critical Difference (CD) test at P = 0.05.

Results and dissection

Effect of biochar on emergence percentage

The effects of biochar on emergence percentage were found not significant as shown in Table (2). It seems that biochar has no effects on the emergence percentage of peanuts specifically, which agreed

with [23]. However, a study showed that the emergence percentage of *Robinia pseudoacacia* L. seeds in biochar-treated soils reached a peak of 2 to 3 days faster than the control [24]. These results were compatible with a study done by [25] that observed the emergence percentage of tomato seeds show 2 to 3 days faster with biochar-amended soil (data not published). Many studies have found no harmful effects of biochar on seed germinations or emergence percentage even with its nanoparticles [23].

Effect of biochar on peanut growth parameters

The growth indices of peanuts were collected two times (40 and 80 days after planting). Results are shown in Tables 2 and 3. Plant height was significantly affected by biochar addition at both measurements. The greatest increase in plant height relative to control was 58 % and 63 % with the addition of 5 % B + 10 t ha⁻¹ GM after 40 and 80 days respectively. The number of branches was significantly affected by biochar addition as well, with the maximum branches noticed at 5 % B + 10 t ha⁻¹ GM. On the other hand, no significant increase was seen in the number of leaves indicator. This may reflect an improvement

in root morphology [26][27], or increase in nutrient retention and water-holding capacity [28].

Table 2. Effect of biochar and organic manure on growth parameter of peanuts after 40 DAS

| Treatment | Emergence percentage | Plant height | Number of branches | Number of leaves |
|----------------------------------|----------------------|---------------------|--------------------|------------------|
| Control | 4.000 | 14.333 ^d | 3.000 ^b | 4.000 |
| 10 t ha ⁻¹ GM | 3.666 | 18.666 ^c | 3.333 ^b | 3.666 |
| 2 % B + 10 t ha ⁻¹ GM | 4.000 | 20.333 ^b | 3.666 ^b | 4.000 |
| 5 % B + 10 t ha ⁻¹ GM | 4.333 | 22.666 ^a | 4.666 ^a | 4.333 |
| F-test | NS | S | S | NS |
| CV % | 26.676 | 3.167 | 10.169 | 24.653 |
| CD (P = 0.05) | - | 1.206 | 0.745 | - |

NS = nonsignificant, S = significant, CV = coefficient of variation and CD = Critical Difference

Table 3. Effect of biochar and organic manure on growth parameter of peanuts after 80 DAS

| Treatment | Plant height | Number of branches | Number of leaves | Number of flowers |
|----------------------------------|---------------------|---------------------|---------------------|--------------------|
| Control | 16.333 ^d | 4.333 ^c | 14.666 ^d | 1.666 ^c |
| 10 t ha ⁻¹ GM | 21.333 ^c | 5.666 ^b | 18.333 ^c | 2.333 ^c |
| 2 % B + 10 t ha ⁻¹ GM | 24.333 ^b | 6.333 ^{ab} | 20.333 ^b | 4.333 ^b |
| 5 % B + 10 t ha ⁻¹ GM | 26.666 ^a | 7.000 ^a | 23.666 ^a | 6.666 ^a |
| F-test | S | S | S | S |
| CV % | 3.445 | 6.387 | 2.296 | 16.026 |
| CD (P = 0.05) | 1.529 | 0.745 | 0.880 | 1.206 |

A number of branches were significantly affected by biochar in both measurements (after 40 and 80 DAS) as shown in Tables 2 and 3. Treatment of 5 % B + 10 t ha⁻¹ GM gave the uppermost number of branches and considerably varied from control and manure treatments. Many studies have also reported that biochar affected plant growth in particular the number of branches [29][30][31].

A number of leaves were not affected by biochar addition in the first measurement (after 40 DAS), but the effect was significant at the second measurement (after 80 DAS). 5 % B + 10 t ha⁻¹ GM gave the highest number of leaves and differ significantly from other treatments. This enhancement in plant growth was also reported by [32][33][30][34]

The number of flowers was high in a 5 % B + 10 t ha⁻¹ GM treatment and statistically differ from other treatments. 2 % B + 10 t ha⁻¹ GM treatment had also changed the number of flowers and gave the second count of flowers and statistically differ from control and manure treatments Table (3). It is noticeable that biochar could enhance plant flowering as a result increase plant yield. Many other studies have reported that biochar had an increase in plant growth and flowering [29][34][35].

Effects of biochar on Peanut yield parameters

The number of pods was significantly affected by biochar Table 4. The addition of 2 % and 5 % biochar increased peanut yield by 50 % and 82 % respectively compared to the control. Without biochar, goat manure increased peanut yield by 20 %. [35] reported that 10 t ha⁻¹ biochar application combined with organic fertilizer in infertile soil increased peanut yield by 50 %. Similarly, the biochar application rate of 10 t ha⁻¹ significantly increased peanut pod yield by 23 % compared to the inorganic fertilizer treatment [36]

Table 4. Effect of biochar and organic manure on yield attribute,

yield of peanut crop

| Treatment | Number of pods | Weight of pods q/h | Straw yield q/h | Harvest index |
|----------------------------------|---------------------|--------------------|--------------------|---------------------|
| Control | 9.666 ^d | 11.41 ^d | 30.90 ^d | 37.00 ^c |
| 10 t ha ⁻¹ GM | 19.000 ^c | 14.29 ^c | 34.50 ^c | 41.43 ^b |
| 2 % B + 10 t ha ⁻¹ GM | 22.333 ^b | 17.12 ^b | 39.60 ^b | 43.23 ^{ab} |
| 5 % B + 10 t ha ⁻¹ GM | 26.666 ^a | 20.80 ^a | 45.90 ^a | 45.46 ^a |
| F-test | S | S | S | S |
| CV % | 7.237 | 2.737 | 2.783 | 3.597 |
| CD (P = 0.05) | 2.808 | 0.865 | 2.098 | 2.994 |

The weight of pods as shown in Table 4, was significantly affected by biochar. As seen with the number of leaves, number of branches, and the number of flowers, 5 % B + 10 t ha⁻¹ GM gave the greatest number of pods. The result is in agreement with [35][37][30].

The straw yield was also affected by biochar (Table 4). The addition of 5 % B + 10 t ha⁻¹ GM cause a 50 % increase in straw yield compared to the control. Biochar enhanced most of the growth and yield parameters measured which was also noticed by other studies [36][38][31][39][22][27].

The harvest index was high in 5 % B + 10 t ha⁻¹ GM treatment and vary significantly from other treatments Table 3. Adding less biochar (2 %) would not increase harvest much when 10 t ha⁻¹ GM was added. In general, biochar improved plant yield and production here and elsewhere [32][37].

Conclusion

The objective of this study was to evaluate the effects of biochar in combination with goat manure on peanut yield in sandy soil. Results showed that biochar had the potential to improve peanut yield when organic fertilizer was practiced for growers. Our study conclude that a high dose of biochar (5%) gave the highest peanut yield which increased the yield by 82 %. In connection with this study, many studies reported that biochar increased plant productivity, with an average yield increase between 10 % to 42 % [40]. The following study should determine if applying more levels of biochar (for example 10 % and 20 %) would increase peanut yield.

Acknowledgements

The authors are thankful to the Agriculture Collage of Sebha University for supporting this work and Authors thank anonymous reviewers for their valuable comments and constructive suggestions towards substantial improvement of this article and the organizers of the sixth international conference of Sebha University.

REFERENCE

- [1]- Bogino, P., Banchio, E., Rinaudi, L., Cerioni, G., Bonfiglio, C., and Giordano, W. (2006). Peanut (*Arachis hypogaea*) response to inoculation with *Bradyrhizobium* sp. in soils of Argentina. *Annals of applied biology*, **148**(3):207-212.
- [2]- Giller, K. E. (2001). *Nitrogen fixation in tropical cropping systems*. CAB Publishing, Wallingford, UK.
- [3]- USDA, (2012) *United States Department of Agriculture National Agricultural Statistics Service*, Washington.
- [4]- Joseph, S., and Lehmann, J. (2009). *Biochar for environmental management: science and technology* (no. 631.422 B615bi). London, GB: Earthscan.
- [5]- Woolf, D., Amonette, J. E., Street-Perrott, F. A., Lehmann, J., and Joseph, S. (2010). Sustainable biochar to mitigate global climate change. *Nature communications*, **1**(1): 56-64.
- [6]- Kuzyakov, Y., Bogomolova, I., and Glaser, B. (2014). Biochar stability in soil: decomposition during eight years and transformation as assessed by compound-specific 14C analysis. *Soil Biology and Biochemistry*, **70**: 229-236.

- [7]- Glaser, B., Lehmann, J., Steiner, C., Nehls, T., Yousaf, M., and Zech, W. (2002). Potential of pyrolyzed organic matter in soil amelioration. In 12th ISCO conference'. Beijing **421**: p. 427.
- [8]- Thies, J. E., and Rillig, M. C. (2012). *Characteristics of biochar: biological properties*. In *Biochar for environmental management* pp. 117-138. Routledge.
- [9]- Novak, J. M., Busscher, W. J., Watts, D. W., Amonette, J. E., Ippolito, J. A., Lima, I. M., and Schomberg, H. (2012). Biochars impact on soil-moisture storage in an ultisol and two aridisols. *Soil Science*, **177**(5): 310-320.
- [10]- Hass, A., Gonzalez, J. M., Lima, I. M., Godwin, H. W., Halvorson, J. J., and Boyer, D. G. (2012). Chicken manure biochar as liming and nutrient source for acid Appalachian soil. *Journal of Environmental Quality*, **41**(4): 1096-1106.
- [11]- Lentz, R. D., and Ippolito, J. A. (2012). Biochar and manure affect calcareous soil and corn silage nutrient concentrations and uptake. *Journal of environmental quality*, **41**(4): 1033-1043.
- [12]- Uzoma, K. C., Inoue, M., Andry, H., Fujimaki, H., Zahoor, A., and Nishihara, E. (2011). Effect of cow manure biochar on maize productivity under sandy soil condition. *Soil use and management*, **27**(2): 205-212.
- [13]- Kookana, R. S., Sarmah, A. K., Van Zwieten, L., Krull, E., and Singh, B. (2011). Biochar application to soil: agronomic and environmental benefits and unintended consequences. *Advances in agronomy*, **112**: 103-143.
- [14]- Lehmann, J., Rillig, M. C., Thies, J., Masiello, C. A., Hockaday, W. C., and Crowley, D. (2011). Biochar effects on soil biota—a review. *Soil biology and biochemistry*, **43**(9): 1812-1836.
- [15]- Spokas, K. A., Cantrell, K. B., Novak, J. M., Archer, D. W., Ippolito, J. A., Collins, H. P., and Nichols, K. A. (2012). Biochar: a synthesis of its agronomic impact beyond carbon sequestration. *Journal of environmental quality*, **41**(4): 973-989.
- [16]- Jeffery, S., Verheijen, F. G., van der Velde, M., and Bastos, A. C. (2011). A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis. *Agriculture, ecosystems and environment*, **144**(1): 175-187.
- [17]- Sohi, S. P., Krull, E., Lopez-Capel, E., and Bol, R. (2010). A review of biochar and its use and function in soil. *Advances in agronomy*, **105**: 47-82.
- [18]- Glaser, B., and Birk, J. J. (2012). State of the scientific knowledge on properties and genesis of Anthropogenic Dark Earths in Central Amazonia (terra preta de Índio). *Geochimica et Cosmochimica acta*, **82**: 39-51.
- [19]- Atkinson, C. J., Fitzgerald, J. D., and Higgs, N. A. (2010). Potential mechanisms for achieving agricultural benefits from biochar application to temperate soils: a review. *Plant and soil*, **337**: 1-18.
- [20]- Elzobair, K. A., Stromberger, M. E., Ippolito, J. A., and Lentz, R. D. (2016). Contrasting effects of biochar versus manure on soil microbial communities and enzyme activities in an Aridisols. *Chemosphere*, **142**: 145-152
- [21]- Major, J., Rondon, M., Molina, D., Riha, S. J., and Lehmann, J. (2010). Maize yield and nutrition during 4 years after biochar application to a Colombian savanna oxisol. *Plant and soil*, **333**: 117-128.
- [22]- Liu, X., Zhang, A., Ji, C., Joseph, S., Bian, R., Li, L., and Paz-Ferreiro, J. (2013). Biochar's effect on crop productivity and the dependence on experimental conditions—a meta-analysis of literature data. *Plant and soil*, **373**: 583-594.
- [23]- Zhang, K., Wang, Y., Mao, J., and Chen, B. (2020). Effects of biochar nanoparticles on seed germination and seedling growth. *Environmental pollution*, **256**: 113409.
- [24]- Bu, X., Ji, H., Ma, W., Mu, C., Xian, T., Zhou, Z., and Xue, J. (2022). Effects of biochar as a peat-based substrate component on morphological, photosynthetic and biochemical characteristics of *Rhododendron delavayi* Franch. *Scientia Horticulture*, **302**: 111148.
- [25]- Elzobair, K. A. (2017) A field study on the tomato crop in Wadi Ataba (Unpublished work).
- [26]- Berek, A. K., and Hue, N. V. (2013). Improving soil productivity with biochars. *Annual report, ICGAI, Yogyakarta, Indonesia*, 1-23.
- [27]- Zheng, J., Wang, S., Wang, R., Chen, Y., Siddique, K. H., Xia, G., and Chi, D. (2021). Ameliorative roles of biochar-based fertilizer on morpho-physiological traits, nutrient uptake and yield in peanut (*Arachis hypogaea* L.) under water stress. *Agricultural Water Management*, **257**: 107129.
- [28]- Wang, B., Gao, B., Zimmerman, A. R., Zheng, Y., and Lyu, H. (2018). Novel biochar-impregnated calcium alginate beads with improved water holding and nutrient retention properties. *Journal of environmental management*, **209**: 105-111.
- [29]- Alwan, S. H., Owen, M. A., Omar, O. A., and Salih, Z. K. (2023). Effect of Biochar Application and Foliar Application of Gibberellin and Cytokinin on Growth and Flowering of Chinese Carnations *Dianthus chinensis*. In IOP Conference Series: *Earth and Environmental Science* **1158**(4): p. 042016). IOP Publishing.
- [30]- Garamu, T. (2019). Effect of different source and rates of biochar application on the yield and yield components of mungbean on the acidic soil in western Ethiopia. *Academic Research Journal of Agricultural Science and Research*, **7**(7): 530-538.
- [31]- Wisnubroto, E. I., Utomo, W. H., and Indrayatie, E. R. (2017). Residual effect of biochar on growth and yield of red chili (*Capsicum annum* L.). *Journal of Advanced Agricultural Technologies*, **4**(1), 28-31.
- [32]- Rahayu, M., Nurmallasari, A. I., and Aini, N. N. (2022, April). Effect of various types and doses of biochar on hybrid maize growth. In IOP Conference Series: *Earth and Environmental Science* **1016**(1): p. 012053). IOP Publishing.
- [33]- Eche, C. O., and Okafor, O. E. (2020). Control potential of some indigenous biochars against *Meloidogyne incognita* [(Kofoid and White) Chitwood] in tomato (*Solanum lycopersicum* L.). *Journal of Entomology and Nematology*, **12**(2): 32-38.
- [34]- Conversa, G., Bonasia, A., Lazzizzera, C., and Elia, A. (2015). Influence of biochar, mycorrhizal inoculation, and fertilizer rate on growth and flowering of *Pelargonium* (*Pelargonium zonale* L.) plants. *Frontiers in plant science*, **6**: 429.
- [35]- Elangovan, R. and Sekaran, N. Chandra (2014). Effect of biochar application on growth, yield and soil fertility status in cotton. *Asian Journal of Soil Science*, **9**(1): 41-49.
- [36]- Wu, P., Ata-Ul-Karim, S. T., Singh, B. P., Wang, H., Wu, T., Liu, C., and Chen, W. (2019). A scient metric review of biochar research in the past 20 years (1998–2018). *Biochar*, **1**: 23-43.
- [37]- Xu, C. Y., Hosseini-Bai, S., Hao, Y., Rachaputi, R. C., Wang, H., Xu, Z., and Wallace, H. (2015). Effect of biochar amendment on yield and photosynthesis of peanut on two types of soils. *Environmental Science and Pollution Research*, **22**: 6112-6125.
- [38]- Tian, X., Li, C., Zhang, M., Wan, Y., Xie, Z., Chen, B., and Li, W. (2018). Biochar derived from corn straw affected availability and distribution of soil nutrients and cotton yield. *Plos One*, **13**(1): 1-19.
- [39]- Ngulube, M., Mweetwa, A. M., Phiri, E., Njoroge, S. M. C., Chalwe, H., Shitumbanuma, V., and Brandenburg, R. L. (2018).

Effects of biochar and gypsum soil amendments on groundnut (*Arachis hypogaea* L.) dry matter yield and selected soil properties under water stress. *African Journal of Agricultural Research*, **13(21)**: 1080-1090.

- [40]- Joseph, S., Cowie, A. L., Van Zwieten, L., Bolan, N., Budai, A., Buss, W., and Lehmann, J. (2021). How biochar works, and when it doesn't: A review of mechanisms controlling soil and plant responses to biochar. *GCB Bioenergy*, **13(11)**: 1731-1764.